

Unveiling the Physics Behind Extreme AGN Variability at UVI on Jul 12, 2017

Discovery of a “dying” AGN in Arp 187

Ichikawa et al. 2016, PASJ, 68, 9

Ichikawa & Tazaki, ApJ in press.

(arXiv:1706.03071)

Kohei Ichikawa (市川幸平)

JSPS Postdoctoral fellow at Columbia/NAOJ

Collaborators

J. Ueda, M. Shidatsu, T. Kawamuro, K. Matsuoka, H-J, Bae,
R. Tazaki

Unveiling the Physics Behind Extreme AGN Variability at UVI on Jul 12, 2017

Discovery of a “dying” AGN in Arp 187

Ichikawa et al. 2016, PASJ, 68, 9

Ichikawa & Tazaki, ApJ in press.

(arXiv:1706.03071)

“Extreme” luminosity decline in 10^{3-4} yrs

Kohei Ichikawa (市川幸平)

JSPS Postdoctoral fellow at Columbia/NAOJ

Collaborators

J. Ueda, M. Shidatsu, T. Kawamuro, K. Matsuoka, H-J, Bae,
R. Tazaki

Key question: AGN lifetime

AGN life time: observationally poorly constrained

- ☑ Total Accretion time scale: 10^{7-8} yr (e.g., Marconi+04)
- ☑ One AGN lifecycle: 10^5 yrs?

(AGN *flickering*; Hopkins & Quataert '10; Novak+11; Schawinski+15;
see also AGN *thermal echo*: Schirmer+16; Kawamuro+17)

How fast do AGN drop its luminosity?

- ☑ 2-3 order of mag in 10^5 yr?

(Hanny's voorwerps e.g., Schawinski+10; Keel+12,15,16, see also Sartori's talk)

- ☑ 1 order of mag in 10 yr

(changing-look QSO; Tohlin & Osterbrock '76 ;LaMassa+15, Ruan+16 etc.)

It might be highly related to AGN accretion disk state?


and/or AGN torus environment?

Here I report additional case of extreme variability

Arp 187

Arp187: Merger remnant LIRG (Rothberg & Joseph '04)

**Good candidate of
AGN feedback study**




ALMA observations (Ueda+14)

- ☑ CO(1-0) associates with host galaxy K band emission
- ☑ 3mm continuum does not associate with CO(1-0) distributions
- ☑ Some dusty outflow occurs? or radio-jet?

Arp 187: VLA observations

VLA 5, 8.5 GHz show bimodal jet lobes




Jet properties

- ✓ Both lobes show spec index $\alpha = -1 \Rightarrow$ synchrotron jet
 - ✓ jet kinetic age: 6×10^4 yr (jet angle = 90° w/ $v = 0.1c$)
 - ✓ **jet core seems absent ($f_{5\text{GHz}} < 210 \mu\text{Jy}$)**
- $\Rightarrow$ current AGN activity might be silent?**

Optical spectrum = NLR activity ($\sim$ kpc)

BPT diagram: NLR (probably) exists




Optical properties: suggest Seyfert class luminosity

- ✓ detections of [OI]6300, H α , [NII], [SII], [OIII], H β
- ✓ [NII]/H α = 0.8, located at composite or AGN
- ✓ $\log L_{[\text{OIII}]} = 42.5$, $\Rightarrow \log (L_{2-10\text{keV}} / \text{erg s}^{-1}) = 44.3$ (e.g., Ueda, KI+15)

mid-IR = torus activity (<10 pc)

Spitzer/IRS (w/4.5''): SB dominant spectra

=> AGN torus emission (<10pc) is faint



$$\begin{aligned} \checkmark L_{12\mu\text{m}} &< 1.5 \times 10^{42} \text{ erg s}^{-1} \Rightarrow L_{2-10\text{keV}} < 2.8 \times 10^{42} \text{ erg s}^{-1} \\ &\Rightarrow L_{14-195\text{keV}} < 5.9 \times 10^{42} \text{ erg s}^{-1} \end{aligned}$$

(see also Gandhi+09; Asmus+15; Ichikawa+12, +17a)

X-ray (= electron corona $\ll 0.1$ pc)

No previous deep X-ray observations for this target

Swift/BAT all-sky survey (E=14-195 keV)

☑ Not listed in the catalog (Baumgartner+13)

=> $L_{14-195\text{keV}} < 4.7 \times 10^{43} \text{ erg s}^{-1}$

=> consistent w/IR upper limit: $L_{14-195\text{keV}} < 5.9 \times 10^{42} \text{ erg s}^{-1}$

ROSAT all-sky survey (E=0.5-2 keV)

☑ Not listed in the catalog (Voges+99)

=> $L_{0.5-2\text{keV}} < 8.8 \times 10^{42} \text{ erg s}^{-1}$

=> consistent w/IR upper limit: $L_{0.5-2\text{keV}} < 2.0 \times 10^{42} \text{ erg s}^{-1}$ ($\Gamma=1.9$; e.g., Brightman+13)

Not located in the XMM-slew survey region

Suzaku obs. (AO-10) was awarded, but not completed

=> Obs. w/ Nustar, ASTRO-H2, and/or Chandra is crucial

BH mass of Arp 187

Using $M_{\text{BH}} - M_{\text{bulge}}$ relation, we estimate M_{BH} .

☑ $L_{\text{Kband}} \Rightarrow M_{\text{stellar}} = 1.3 \times 10^{11} M_{\text{sun}} \text{ w/n}_{\text{sersic}} \sim 4$ (Rothberg & Joseph '04)

$\Rightarrow M_{\text{stellar}} \sim M_{\text{bulge}}$

$\Rightarrow \mathbf{M_{BH} \sim 7 \times 10^8 M_{\text{sun}}}$ (Kormendy & Ho 2013)

Upper limit of λ_E :


☑ $L_E = 8.4 \times 10^{46} \text{ erg s}^{-1}, L_{\text{bol}} < 8.4 \times 10^{43} \text{ erg s}^{-1}$

$\Rightarrow \mathbf{\lambda_E < 10^{-3}}$

Disk is now ADAF state (Ichimaru '77, Narayan & Yi '94)

BH fundamental plane

If you have either two values of $L_{5\text{GHz (core)}}$, $L_{2-10\text{keV}}$, M_{BH} , then you can estimate the final one.



Merloni+03
Heinz+03
Yuan+14

Upper limit of L_{R} , and M_{BH} are given, then


☑ $L_{2-10\text{keV}} < 4 \times 10^{39} \text{ erg s}^{-1}$ (see Gultekin+09; Yuan & Cui '05)

What is happening in Arp 187?

- ☑ Signs of NLR, jet ($> \text{kpc}$)
- ☑ Absence of torus, radio core ($< 10 \text{ pc}$ scale)

FYI; torus cools down
in $< 100 \text{ yr}$ (KI & Tazaki '17)


AGN “was” active $\sim 10^4 \text{ yr}$ ago,
but currently AGN “is” dead.



“AGN light echo” is powerful tool to know AGN activity history


see Hanny’s Voorwerp (10^{4-5} yr ago; Lintott+09, Schawinski+10, 15; Keel+12)
or the galactic center (500 yr ago; Ponti+10; Capelli+12; Ryu+13; Nakashima+14)
or Fermi bubbles study (10^6 yr ago; Zubovas+11, 12; SU & Finkbeiner '12)

AGN Luminosity drops of Arp 187



L_{AGN} has decreased at least 10^{3-4} times in $\sim 10^{3-4}$ yr

AGN Luminosity drops of Arp 187



L_{AGN} has decreased at least 10^{3-4} times in $\sim 10^{3-4}$ yr

Summary: witnessing the death of AGN

Arp 187 is a LIRG w/ interesting properties

☑ NLR (w/ age $\sim 10^{3-4}$ yr), jet lobe (w/ age $\sim 10^5$ yr) exist

☑ torus, jet core emission is absent


=> AGN “was” active but “is” dead

L_{AGN} : go downs at least 10^{3-4} times in $\sim 10^{3-4}$ yr

**Sudden ($\sim 10^{3-4}$ yr) Death of AGN in Arp 187
with “extreme” L decline (3-4 orders)**

Appendix

Torus cools down in 10-100 yrs



dust clump
energy Eq.

$$\frac{dE_{c,d}}{dt} = -4\pi R_c^2 \langle Q_c \rangle_T \sigma_{\text{SB}} T^4 + \int_{V_{\text{clump}}} n_g \Gamma_{g-d} dV \quad (13)$$

Torus cools down in 10-100 yrs




Figure 2. Time evolution of dust torus SED around dying AGN whose initial luminosity before the quenching is $10^{45} \text{ erg s}^{-1}$. The dashed line represents the SED under the radiative equilibrium temperature structure.